VOLUME 64, NUMBER 1 FEBRUARY 2004

Pacific and American Golden-Plovers: Reflections on Conservation Needs

by Oscar W. Johnson¹

Reprinted with the kind permission of both the author, O.W. Johnson, and the editor of Wader Study Group Bulletin, Humphrey Sitters. This article originally appeared in the Wader Study Group Bulletin 100: (April 2003) pp10–13.

Much has been learned about the biology of Pacific and American Golden-Plovers (*Pluvialis fulva* and *P. dominica*) in recent years. However, most of this information does not address conservation-related questions affecting the future of these birds. We lack accurate demographic figures (guesstimates of world population vary more than 30-fold in each species), and population trends are uncertain. Reports of large spring aggregations (*fulva* in northeastern Mongolia and *dominica* in west-central Indiana) suggest stopover sites of hemispheric importance

that urgently need further evaluation and possible protection. Little is known concerning various anthropogenic threats including contaminants, loss of habitat, and hunting. In contrast to fulva, dominica do not readily co-exist with humans during the nonbreeding season; also much of their South American winter range has been converted to croplands. Thus, of the two species, dominica appears to be the most at risk.

INTRODUCTION

Studying long-distance migrant shorebirds offers many satisfactions and rewards. Not the least of these is that such projects often require the researchers to be migratory as well and follow their subjects to interesting places. For the past couple of decades, my colleagues and I have marked, with unique band combinations, Pacific Golden-Plovers Pluvialis fulva in Hawaii and Alaska, and American Golden-Plovers P. dominica in Alaska. Long-term cooperation from the birds (they are very site-faithful from year-to-year), together with similar data gathered by investigators elsewhere, have enabled us to shed light on various basics including site-fidelity, territorial behaviours, nesting, and longevity (Johnson et al. 2001a,b; Plate 1). While this is valuable information, there are also more far-reaching and difficult questions in the realm of conservation that need investigation. As wader biologists well know, the future of all longdistance migrants rests on such factors as habitat preservation and improved knowledge of population trends. In the following commentary, my intent is to call attention to conservationrelated matters that bear on the well-being of Pacific and American Golden-Plovers - especially voids in knowledge and threats of anthropogenic origin.




Plate 1. In our studies of Pacific Golden-Plovers on O'ahu wintering grounds (Johnson *et al.* 2001b), the life expectancy of most birds after banding was around 5 years, but there were exceptional individuals. This "participant" in the research lived to a minimum age of 21 years 3 months, a new world record for the species (for details, see Johnson *et al.* 2003).

WORLD POPULATIONS AND TRENDS

At present we have nothing better than educated guesses as to how many of these plovers actually exist, and almost no information on population trends. Estimates of population size are far from consensus: fulva, 125,000 (including 16,000 breeding in North America where the species nests only in Alaska) to 2.6 million breeding pairs; dominica, 150,000+ to 2.5 million breeding pairs. The lower estimates were made by Morrison et al. (2001b), the higher by Byrkjedal & Thompson (1998). Morrison et al. consider their figures to be of low accuracy, but probably "in the right order of magnitude"; the high-end estimates are likely too optimistic. Monitoring trials in Alaska during the 2002 breeding season suggest that the actual statewide population of fulva exceeds the estimated 16,000 birds (C. Wightman, pers. comm.). The most recent estimate for fulva incorporates the Morrison et al. figure of 16,000 nesting in Alaska and places world population at 166,000-216,000 (Delany & Scott 2002). Given the difficulties inherent in counting migrants that range over vast areas of the world, our lack of knowledge concerning their numbers is understandable. Securing the data for improved estimates of global fulva and dominica populations remains an ongoing challenge.

Hawaii contains some of the most important *fulva* wintering grounds in the world, and assessment of plover numbers there continued on page 3

Program Meeting - February 16th

Wildlife biologist Sharon Reilly will speak on Ducks Unlimited's conservation projects in Hawai'i. Through its newest conservation initiative, Wetlands Hawai'i, Ducks Unlimited is working in cooperation with a number of public and private groups to restore, protect, and manage Hawai'i's wetlands. Partners include Ducks Unlimited, the U.S. Fish and Wildlife Service, Natural Resour

Projects in Hawai'i include: using satellite technology to map existing wetlands; in Pearl Harbor working to restore a tidal marsh for Hawaiian stilts; on Kaua'i, working with the Fish and Wildlife Service to restore wetlands and provide lands for taro growers on Hanalei National Wildlife Refuge. One of their biggest efforts is helping with restoration and management planning on the new Kealia National Wildlife Refuge on Maui.

Program meetings are held at Henry Hall Room 109 on the Chaminade University campus, 3140 Wai 'alae Avenue, Kaimuki. Meetings are from 7:30 to 9:30pm. Refreshments are served, and HAS products and publications are available for purchase.

Results of Board Election

By a plurality of votes, the following were elected or re-elected to the Hawaii Audubon Society Board of Directors:

President, Larry Kimmel
Vice President, Liz Kumabe
Treasurer, Sal Pagliaro
Recording Secretary, Tonnie Casey
Director, Carol Bebb
Directors, Chad Castle and Alice Roberts

Directors whose terms were not up for re-election, Arlene Buchholz, Phil Bruner, Jennifer Crummer, and Ron Walker complete our Board.

CORRECTION

The reference to a Red-crested Cardinal at the end of paragraph three in the article "Birding on Maui and Online" by Satoko K. Lincoln should have been a reference to a Northern Cardinal.

'ELEPAIO

ISSN 0013-6069

Managing Editor: Linda Shapin Scientific Editor: Ron Walker

850 Richards Street, Suite 505 Honolulu, HI 96813

Tel: (808) 528-1432 Fax: (808) 537-5294

E-mail: hiaudsoc@pixi.com Website: www.hawaiiaudubon.com

The 'Elepaio is printed on recycled paper and published nine times per year: February, March, April, May, June/July, August/September, October, November, and December/January

Field Trips for 2004

All trips with an * are still in the process of being planned. Details will be provided as the scheduled dates get closer. A donation of \$2 per participant on all field trips is appreciated. Field Trip information is also available on the HAS office answering machine (528-1432) and on our website, <www.hawaiiaudubon.com>

Saturday, March 27th, 8:30 am Kahuku Dunes shoreline walk with Dr. Phil Bruner. A half day or all day trip depending on each individuals available time. We will meet at 8:30 am at the parking lot of Kahuku Golf Course and hike from there along the shoreline to Turtle Bay Resort. We will take the city bus back to Kahuku at the conclusion of the trip (bring \$2 bus fare). On the hike we will see and talk about seabirds, migratory shorebirds, dune formation, coastal plants and their unique adaptations. We will also observe the conservaion concerns in this environment. People should bring plenty of water, a lunch, sunscreen, hat and binoculars. Copies of Hawaii's Birds by Hawaii Audubon Society and Hawaiian Coastal Plants by Mark Merlin are suggested to bring along. This hike is in one of the few remaining undeveloped coastal lands on O'ahu. Call the HAS office for starting location and to register – 528-1432.

Saturdays, April 3rd and 17th, both 7:30am Shorebird Farewell at Paiko Lagoon. A chance to bid our shorebirds (Pacific Golden Plover, Ruddy Turnstone, Sanderling, etc.) goodbye, as they will be leaving for Alaska shortly. Wear old tennis shoes or reefwalkers, and bring sunscreen, water, and lunch. We will meet at Paiko Lagoon. Call Alice to register, 538-3255.

Kolea Watch III - Spring 2004

Kolea Watch III is being planned for spring 2004. In collaboration with the Hawai'i Nature Center & the U.S. Fish and Wildlife Service, Dr. Wally Johnson (Montana State University) and colleagues will attach temporary radio transmitters to Kolea and then monitor their migration. Similar research over the past two spring seasons produced important new information about the destinations of Hawai'i's plovers in Alaska. A paper detailing these findings is currently being reviewed for publication in a major ornithological journal (Wilson Bulletin). The manuscript includes acknowledgments to Hawaii Audubon Society and Kolea Watch donors for their invaluable support of the project.

If you want to help the investigators with their continuing studies, please consider an "adopt-a-plover" donation. Each transmitter costs \$135, but assistance in any amount will be most welcome. Anyone buying a transmitter will have "adopted" the bird that carries it to Alaska, and will be informed if/when/where "their" bird is found. Or, maybe you would consider helping to fund aerial monitoring in Alaska. The costs there are about \$100 per hour of flight time, and the average survey takes about 5 hours. Donations for Kolea Watch can be sent to: Hawaii Audubon Society, 850 Richards Street, Honolulu, HI, 96813-4709. Checks should be made payable to Hawaii Audubon Society and earmarked "Kolea Fund."

Update on Studies of the Mid-Pacific Migration of Kolea Between Hawai'i and Alaska

Over the past three spring seasons (2001-2003), Dr. Wally Johnson, Montana State University, radio-tagged a total of 55 Pacific Golden-Plovers(Pluvialis fulva) at wintering ground study sites on O'ahu. Following their northward migration (most birds deserted winter territories in late April), 15 of them were found in Alaska during aerial surveys by cooperating U.S. Fish and Wildlife Service biologists. Signals from the birds were heard in three regions: Copper River Delta, King Salmon, and Kotzebue. One individual made the transpacific flight from O'ahu to the Alaska Peninsula in a minimum time of 70 hours at a minimum flight speed of 56 km/hour. Present findings, together with Johnson's earlier records of 45 plovers radio-tagged on O'ahu (13 were detected in Alaska), indicate a major Hawai'i-Alaska migratory connection for this species, and suggest that plovers wintering on O'ahu nest throughout the known Alaska breeding range.

The radio transmitters are temporary in that they fall off the plovers during the nesting period on the northern tundra. Following the breeding season, 84% of the sample birds returned to O'ahu and reoccupied their previous winter territories. Notably, this return rate was identical to returns of marked plovers that had not been radio-tagged indicating that the temporary attachment of a tiny transmitter to feathers on the back caused the birds no harm.

Johnson's studies (the first of their kind anywhere in the central Pacific region) represent a major step forward in understanding trans-Pacific migration between Hawai'i and Alaska. His earlier findings have already been published in ornithological journals, and a paper detailing the last three seasons is currently being reviewed for publication. Portions of the research project have been funded by the Hawaii Audubon Society. Further radio-tagging is being planned for spring 2004 — see the announcement in this issue entitled "Kolea Watch."

continued from page 1

should be of high priority as a baseline for gauging future trends in the mid-Pacific flyway. The most recent figures are 74,000 birds for the main Hawaiian Islands in 1949, and 15,173 for Oahu in 1968 (see Johnson & Connors 1996). Over the ensuing years, urbanization has produced much additional grassland and the state may host more wintering birds now than ever before. We have no meaningful trend information for the species anywhere in the world. Although reports suggest that numbers have declined at the southern end of the winter range in Australia and New Zealand (Wilson 2001), it is uncertain whether this actually indicates fewer birds or merely reflects short-stopping elsewhere in favourable places such as Hawaii.

Limited trend data for *dominica* are in some cases conflicting, but overall suggest that the species is in decline (Johnson & Connors 1996, Gratto-Trevor 1998, Morrison *et al.* 2001a). From 1996 to 2000, J.P. Isacch (*in litt.*) made counts of plovers at Medaland Ranch, which is one of the primary remaining wintering grounds in Argentina. His results were encouraging in that the number of birds found was similar to what Myers and Myers (1979) reported in the same area nearly a quarter of a century ago. However, because Medaland contains well-preserved habitat in a key wintering region, counts there might remain relatively stable despite losses elsewhere. Additional systematic monitoring of *dominica* in South America is an urgent conservation need.

MIGRATORY ROUTES AND STOPOVERS

The routes followed by these plovers (among the longest flights of all shorebirds) are generally understood, but we lack insight concerning movements across remote regions such as *fulva* on trans-Asian, and southwest Pacific routes, or *dominica* on trans-Amazonian pathways. Although stopovers by these birds often appear to be opportunistic, there are indications of important re-fueling sites of which we have only partial knowledge.

Very few fulva occur in the large spring aggregations of shorebirds found along the Asian coast (Barter et al. 1998, 2000, 2002; Moores 1999). Presumably, the birds are overflying the coast in favour of one or more inland routes to breeding grounds in Siberia (Johnson & Connors 1996, Ming & Kraaijeveld 2000). This raises important questions about stopovers during trans-Asian passage. In this regard, Oleg Goroshko's 1993–1996 spring observations (in litt.) of about 50,000 fulva in the Torey Depression of NE Mongolia and adjacent Russia (see www.wetlands.org/RDB/Ramsar_Dir/RussianFed/RU019D02.htm) are of great interest. Goroshko's numbers suggest a site of hemispheric importance for the species, and emphasize the need for further study of fulva in the Torey steppe/lake region.

American Golden-Plovers are well known for their annual migratory pattern, which involves a mid-continent route through North America in the spring and an offshore trans- Atlantic flight in fall (Johnson & Connors 1996). Stopovers by spring migrants are somewhat unpredictable, though major gatherings occur in Indiana and Illinois (Johnson & Connors 1996) where the birds feed on earthworms and insects in agricultural fields. In Indiana, they find untilled soybean fields harvested the previous fall especially attractive (Braile 1999). West-central Indiana (Benton County and parts of White County) appears to be particularly important for the species. In this relatively small area, Braile (1999) estimated that the number of migrant dominica foraging during stopovers ranged from 42,000 to 84,000 - a substantial fraction of the world's total population. Clearly, this locale is of major significance as a possible WHSRN (Western Hemisphere Shorebird Reserve Network) site, and it is imperative that its use by dominica be evaluated further. In a region like Indiana, bird habitat is dependent on agricultural practices that vary with market demands and economic factors. Despite this uncertainty, there may be long-term strategies for managing the land in ways beneficial to farmers and plovers alike.

continued on page 5

Many Thanks!

The Board of Directors and the Staff of the Society wishes to thank those who responded between November 2003 and January 1, 2004 to the 2003 Annual Appeal, and other recent donors. These contributions will be used to continue and expand our many programs that benefit Hawai'i's native wildlife. Many thanks to:

Hawaii Audubon Society members: Karen M. Anderson, Diane Aoki, Bob Asanoma, Maile and John Bay, Brenda Becker, Phillip Bruner, J and M Buchholz, Thomas R. Chell, Barbara L. Coles, Ms. Kathryn Copenhaver, Coleen Cory, Evelyn Cox, Dr. Ronald Crombie, S.F. Cushman, Scott R. Derrickson, Eugene And Rita Donaldson, Susan Dursin, Richard R. Eakin, Michael J. Fischer, Joel Fithian, James H. Fleming, Leonard Freed, Hubert Frings, Lillian Fujii and Steve Hayashi, Norm Garon, J. Richard Gauthey, Thomas Goodier, Kay J. Green, Alice Gruber, Mrs. Ann Guild, John H. Haas, Clifford W. Hand, Dr. Ray and Karen Harris, Emily Herb, Robert Hill, Lynn M. Hodgson, Kamal Islam, Ph.D., Pearl C. Johnson, Dr. Oscar W. Johnson, Wendy B. Johnson, Virginia P. Johnson, Bud Johnson, Kathryn Keaton, Brian Keesling, Dave Kemble, Dr Lawrence J. Kimmel, Warren B. King, Dean Kleinhans, Elizabeth Kumabe, Janelle Kunihiro, Edwin F. Laak, Ron Lau, Mr. And Mrs. Eric Lebuse, Robin L.C. Leong, Diane M. Privette and Keith A. Loll, Cathy Lowder, John J. Mahoney, Ruth Mcangus, John T. Mcbride, Jr., Anne Mckay, Matthew Medeiros, Elizabeth Miller, Daniel Nelson and Craig Gorsuch, Diane J. Nielsen, Bruce G. Noll, Kit & Buzz Nordeen, Mitchell Oliphant, Linda Paul, Dorothy Perry, Jennie L. Phillips, Thane Pratt, Diane E. Quick, Mark J. Rauzon, George Reed, Joby Rohrer, Susan Schenck, Kurt R. Schwarz, Kazuo Senda, Leon Slawecki, Andrew Starrett, Karen And Allan Stockton, Talbert K. Takahama, Lynn R. Thomas, Emilia Thomas, Walter Y. Tokushige, Dr. Rowena Toomey, Phyllis Turnbull, Lee & Melinda Varian, Ann E. Viets, Betty M. Vitousek, Mrs. Lucile Waldron, Ronald L. Walker, Gerald J. Zasa, Robert Zimmerman.

Joint (National and HAS) members: Anne Anderson, Michael and Joy Anderson, Ms Nancy M Bannick, Maile and John Bay, Arlene Buchholz DVM, Lillian T Ching, Ms. Noreen Chun, Cassandra Dekramer, Mary Dobson, Marilyn F Dunlap, John L Earle, William Eddy, Ms. Patricia G. Engelhard, Jim and Elaine Fessenden, June Firing, Tony and Gloria Fraiola, Stephanie Frazier, J. Frierson, Denny Geiler, Scott Gier, Cheryl and John Goody, Bill and Norma Gorst, Richard Gray, Michael G. Hadfield, Mrs Richard H. Hagemeyer, Emmett Hall, James Harrow, Hawaiian Wildlife Tours, Suzie Heidrich, Hike Maui, Robert B. Hill, Brian R. Jenkins, Lorna Jeyte, Annette Kaohelaulii, Diana King, Nancy Koelper, Abigail S. Laros, Robert Leclair, G. Leo, Peter Lindhout, Ana Lokelani, Ms. Karen L. Loomis, Carol Loose, Craig Martin, Dennis Masui, Sharie Matsumoto, Takeshi Motomura, Kenneth Nakagawa, Glynnis Nakai, Akira Ogata, Kyle Ono, Chuck Pucevich, Louise Ripple, Allan and Shirley Samuelson, Frank and Laura Sayre, Gale Scudder, Kazuo Senda, Joanne Shigekane, Kathy Shimata, Edward Shiroma, Peggy Spencer, Carolyn Stephens, Dennis Tataki, Clifford Tervola, James Thropp, Ann Y. Todd, Carol Tomioka, Troy Vought and Mary Taisala, Fay Warshauer, David And Abby Watson, Lorrin Wong, Wendy T. Yamaguchi.


Greenhouse Frog Invasion

Greenhouse frogs and coqui have been spreading in Hawai'i primarily through infested materials in the plant trade. In the case of coqui, the distribution of the frogs in the State is relatively easy to track due to their distinctive vocalization. Greenhouse frogs



are not. The call is softer and noticed less often. Currently, it is unclear what the distribution of the greenhouse frog is on the islands of Hawai'i, Maui, and O'ahu. There are attempts to locally control populations of coqui and greenhouse frogs with citric acid solutions. This work has been conducted by staff of the Hawai'i Dept. of Agriculture, USDA - Wildlife Services, and Hawai'i Dept. of Land and Natural Resources. Key issues with treatment is relative efficacy in relation to environmental conditions and re-invasion of treated sites due high local populations of frogs.

Earl Campbell, Ph.D.
Invasive Species Coordinator
USFWS - Pacific Islands Fish and
Wildlife Office
Honolulu, Hawai'i
(808) 792 - 9400



CONTAMINANTS

P. fulva wintering at various insular Pacific sites demonstrate a remarkable ability to co-exist with humans in urban environments (Johnson & Connors 1996, Stinson et al. 1997, Johnson et al. 2001b. Beichle 2001). The removal of forest and brush on Pacific islands and atolls tends to produce grassy habitats that support large numbers of wintering plovers. While this initially seems favourable, there may be undesirable trade-offs of which we are unaware. Various pesticides are commonly used in the Pacific region often in situations where there are numerous plovers (golf courses in Hawaii, for example), and the effects of these chemicals on winter-resident fulva have yet to be determined. Inter-year plover survival rates at our Oahu study sites are relatively high at about 80% (Johnson et al. 2001b, O.W. Johnson Wader Study Group Bulletin 12 Bulletin 100 April 2003 unpubl.), but this might be misleading since contamination at these sites is probably minimal. Thus, our survival findings may not be broadly applicable either in Hawaii or elsewhere across the insular Pacific.

It seems likely that both species are carrying pesticides, but whether at harmful levels is uncertain. Also, there may be variations from one flyway to another. One of the most hazardous routes might be the corridor traversing mid-continent North American farmlands. Spring sightings indicate that dominica spend several weeks in gradual northward passage (Johnson & Connors 1996), and probably are exposed to numerous agrochemicals at feeding stops along the way. Because of the immense geographic areas over which both species move annually, substantial reduction in hazards from contaminants may be impossible. There are situations, however, in which at least modest improvement might be achieved. For example, I have observed insecticides being applied on wintering grounds (government owned property on Oahu) in April when birds are fattening for spring migration. Since plover departure from Hawaii is predictable in late April, it should be feasible to modify spraying schedules and defer chemical use until a postdeparture date in May. Possibly, such a regulation could be mandated by federal and/or state agencies.

MORTALITY

Unknown numbers of *fulva* are being lost annually to human exploitation (i.e., commercial and subsistence hunting) particularly in SE Asia, the Philippines, and Indonesia (Alonzo-Pasicolan 1990, Tang & Wang 1995, Johnson & Connors 1996, Ma *et al.* 1998). It appears that American Golden-Plovers are still being hunted in Barbados and parts of South America (Johnson & Connors 1996). During a visit to Barbados in fall 2001, S. Gillings (*in litt.*) observed shooting ponds "full of wooden wader decoys" and "aviaries containing live decoys including two American Golden-Plovers". Though the take of *dominica* may not be excessive (Rappole 1995), we have no clear record of the overall numbers being killed.

A potential mortality factor that could become significant in the future has to do with *fulva* at airports. The latter often contain grassy areas occupied by wintering territorial plovers, and their runways attract passing migrants. Collisions with commercial aircraft have occurred in Hawaii usually in the fall, and these typically involve naive juvenile birds on their first southward migration (Linnell et al. 1996, Johnson & Connors 1996). Thus far, the problem has been relatively minor (i.e., small numbers of birds killed and no aircraft accidents), but new quieter jets now coming on line are more apt to take inexperienced birds by surprise. Resident territorial plovers (mostly adults) are habituated to aircraft and skilled at avoiding them; also these birds tend not to frequent runways. By contrast, inexperienced juveniles often travel in flocks likely to alight on runways. It would be unfortunate if officials unfamiliar with these features of fulva were to institute systematic control programs in and around airfields.

Predictably, such efforts will produce vacant habitats (i.e., by removal of territorial individuals) attractive to other plovers from nearby areas, thereby resulting in a lethal sink. The most effective and acceptable management tools at airports are deterrent measures designed to keep fall juveniles away from runways (in some situations, deterring spring migratory aggregations might also be necessary) along with habitat management (Johnson & Connors 1996). Innovative deterrent techniques involving dogs and possibly falcons are likely to be more successful than harassing birds with a vehicle or loud sounds. Habitat management may involve removal or reduction of grassland on some portions of the airport combined with habitat enhancement on others. Although the data might be difficult to obtain, it would be useful to determine what airport policies are in effect in other regions of the fulva winter range. While there are no lethal control programs in Hawaii, such measures at airfields elsewhere could be causing significant plover mortality of which we are unaware.

OTHER COMMENTS

During the nonbreeding season, dominica have an altogether different "personality" than fulva. In contrast to the adaptable nature of fulva in which foraging on residential lawns, roosting on roof-tops, and even accepting hand-outs of food are routine wintering behaviours, dominica remain wary and require pasturelands and coastal wetland habitats in winter (Johnson & Connors 1996, Johnson et al. 2001b, Beichle 2001). The combination of intolerance toward humans plus loss of habitat (much of the pampas region has been converted to cropland unsuitable for dominica), does not bode well for the species. Given their need to forage in short vegetation, any attempts to manage winter range grasslands for the benefit of dominica should incorporate relatively heavy grazing by livestock (see Lanctot et al. 2002). American Golden-Plovers are known to defend territories on wintering grounds in Argentina (Myers & Myers 1979), but how this relates to intra- and inter-year sitefaithfulness is uncertain because these observations did not involve banded individuals. To my knowledge, there have never been studies of marked dominica during the nonbreeding season, and this is a major gap in our understanding of the species.

Fortunately, there are various national and international conservation efforts (familiar to most readers) currently focussed on the preservation of shorebirds. Harrington *et al.* (2002) provide an informative discussion of the new "Program for Regional and International Shorebird Monitoring" (PRISM

continued on page 6

continued from page 5

- see also Skagen et al., this volume), and the "Western Hemisphere Shorebird Reserve Network" (WHSRN) founded in 1986. Collaborative programs like these are vital to the well-being of many shorebird species including fulva and dominica. New findings on population trends, annual breeding success, and threats to habitats (including the potential effects of climate change – see Rehfisch & Crick, this volume) will be especially valuable if they translate into the means necessary to ensure the future of these remarkable birds.

ACKNOWLEDGEMENTS

I am grateful to Humphrey Sitters and Simon Gillings for their helpful reviews of the manuscript.

Department of Ecology, Montana State University, Bozeman, MT 59717 e-mail: owjohnson2105@aol.com

REFERENCES

- Alonzo-Pasicolan, S. 1990. A survey of hunting pressure on waterbirds Lumpur, Malaysia.
- Barter, M.A., D. Tonkinson, J.Z. Lu, S.Y. Zhu, Y. Kong, T.H. Wang, Z.W. Li & X.M. Meng. 1998. Shorebird numbers in the Huang He (Yellow River) Delta during the 1997 northward migration. *Stilt* 33:15–26.
- Barter, M.A., J.R. Wilson, Z.W. Li, Z.G. Dong, Y.G. Cao & L.S. Jiang. 2000. Yalu Jiang National Nature Reserve, north-eastern China – a newly discovered internationally important Yellow Sea site for northward migrating shorebirds. Stilt 37: 14–21.
- Barter, M.A., J.J. Du, H. Wang, Y.Q. Chen, Z.D. Gao, H. Cheng & C.R. Li. 2002. Shorebird numbers in the Yancheng National Nature Reserve during the 2001 northward migration. Stilt 41: 27–34.
- Beichle, U. 2001. Pacific Golden Plovers Pluvialis fulva and other waders on the Samoan Islands: Wintering in a changing Polynesian landscape. Wader Study Group Bull. 96: 40– 46.
- Braile, T.M. 1999. Migration studies of shorebirds in westcentral Indiana. MS thesis, Purdue University, Lafayette, Indiana.
- Byrkjedal, I. & D. Thompson. 1998. Tundra plovers: The Eurasian, Pacific and American Golden Plovers and Grey Plover. T & AD Poyser Ltd., London.
- Delany, S. & D. Scott (compilers). 2002. Waterbird population estimates, 3rd ed. Wetlands International Global Series No. 12, Wageningen, The Netherlands.
- Gratto-Trevor, C.L., V.H. Johnston & S.T. Pepper. 1998. Changes in shorebird and eider abundance in the Rasmussen Lowlands, NWT. Wilson Bull. 110: 316–325.
- Harrington, B.A., S.C. Brown, J. Corven & J. Bart. 2002. Collaborative approaches to the evolution of migration and the development of science-based conservation in shorebirds. Auk 119: 914–921.
- Johnson, O.W. & P.G. Connors. 1996. American Golden-Plover (*Pluvialis dominica*), Pacific Golden-Plover (*Pluvialis*

- fulva). In *The Birds of North America*, No. 201–202 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C.
- Johnson, O.W., P.L. Bruner, A.E. Bruner, P.M. Johnson, R.J. Kienholz & P.A. Brusseau. 2001a. Features of breeding biology in Pacific and American Golden-Plovers nesting on the Seward Peninsula, Alaska. Wader Study Group Bull. 95: 59–65.
- Johnson, O.W., P.L. Bruner, J.J. Rotella, P.M. Johnson & A.E. Bruner. 2001b. Long-term study of apparent survival in Pacific Golden-Plovers at a wintering ground on Oahu, Hawaiian Islands. Auk 118: 342–351.
- Johnson, O.W., P.L. Bruner, P.M. Johnson & A.E. Bruner. in press 2003. A new longevity record for the Pacific Golden-Plover. J. Field Ornithol.
- Lanctot, R.B., D.E. Blanco, R.A. Dias, J.P. Isacch, V.A. Gill, J.B. Almeida, K. Delhey, P.F. Petracci, G.A. Bencke & R.A. Balbueno. 2002. Conservation status of the Buffbreasted Sandpiper: Historic and contemporary distribution and abundance in South America. Wilson Bull. 114: 44–72.
- Linnell, M.A., M.R. Conover & T.J. Ohashi. 1996. Analysis of bird strikes at a tropical airport. J. Wildl. Manage. 60: 935– 945.
- Ma, M., J. Lu, C. Tang, P. Sun & W. Hu. 1998. The contribution of shorebirds to the catches of hunters in the Shanghai area, China during 1997–1998. Stilt 33: 32–36.
- Ming, M. & K. Kraaijeveld. 2000. Recent records of waders in Xinjiang Uygur Autonomous Region, north-western China. Wader Study Group Bull. 92: 25–29.
- Moores, N. 1999. A survey of the distribution and abundance of shorebirds in South Korea during 1998–1999: interim summary. Stilt 34: 18–29.
- Morrison, R.I.G., Y. Aubry, R.W. Butler, G.W. Beyersbergen, G.M. Donaldson, C.L. Gratto-Trevor, P.W. Hicklin, V.H. Johnston & R.K. Ross. 2001a. Declines in North American shorebird populations. Wader Study Group Bull. 94: 34–38.
- Morrison, R.I.G., R.E Gill, Jr., B.A. Harrington, S. Skagen, G.W. Page, C.L. Gratto-Trevor & S.M. Haig. 2001b. Estimates of shorebird populations in North America. Occas. Paper No. 104, Canadian Wildlife Service, Ottawa, Ontario.
- Myers, J.P. & L.P. Myers. 1979. Shorebirds of coastal Buenos Aires Province, Argentina. *Ibis* 121: 186–200.
- **Rappole**, **J.H.** 1995. *The ecology of migrant birds: a neotropical perspective*. Smithsonian Inst. Press, Washington, D.C.
- Stinson, D.W., G.J. Wiles & J.D. Reichel. 1997. Occurrence of migrant shorebirds in the Mariana Islands. J. Field Ornithol. 68: 42–55.
- Tang, S. & T. Wang. 1995. Waterbird hunting in East China. Asian Wetland Bureau Publ. No. 114, Kuala Lumpur, Malaysia.
- Wilson, J.R. 2001. The Australian Wader Studies Group population monitoring project: Where to now? Perspectives from the chair. *Stilt* 39: 13–26.

HAS Products and Publications

May be purchased by mail, on our website www.hawaiiaudubon.com via PayPal, or in person at our office to avoid mailing costs.

Hawai'i's Birds by the Hawaii Audubon Society, 5th edition, 1997. Over 150 color photographs and illustrations. \$12.95 per copy (\$9.95 plus \$3.00 postage and handling).

ON SALE! Voices of Hawaii's Birds by Douglas Pratt and the Hawaii Audubon Society, 1995. Two 60 minute tapes of songs of more than 100 species of birds found in Hawaiii. Includes a booklet. \$8.00 (\$5.00 plus \$3.00 postage and handling).

Hawai'i's Rare & Endangered Birds Notecards with artwork by renowned Island artist Patrick Ching. Eight greeting cards with envelopes. \$9.50 per box (7.00 plus 2.50 postage and handling).

Treasures of O'ahu Map. A fun and informative self-guided tour of O'ahu including hiking, birding, and ancient cultural sites. \$5.00 (\$4.00 plus \$1.00 postage and handling).

The Edge of Forever The Society's 60th Anniversary Commemorative Poster by Richard Pettit. Now only \$10.00 (\$5.00 plus \$5.00 postage, mailing tube, and handling).

NEW! HAS 2004 one-page calendar featuring a lovely Tropicbird photo by T. Dove. \$4.00 (\$3.00 plus \$1.00 postage and handling).

'Elepaio logo patch \$3.75 (\$3.00 plus \$.75 postage and handling).

Checklist of the Birds of Hawai'i—2002 by R.L. Pyle. Lists all taxa naturally occurring in Hawai'i and introduced species that have established viable populations. \$3.50 (\$3.00 plus \$.50 postage and handling).


Checklist of the Birds of the Mariana Islands by James D. Reichel and Philip O. Glass, 1991. Lists all taxa naturally occurring in the Marianas and introduced species that have established viable populations. \$3.00 (\$2.00 plus \$1.00 postage and handling).

Checklist of the Birds of Micronesia by P. Pyle and J. Engbring, 1985. Lists all taxa naturally occurring in Micronesia and introduced species that have established viable populations. \$3.00 (\$2.00 plus \$1.00 postage and handling).

Field Card of the Birds of Hawai'i by R. L. Pyle and R. David, 1996. A pocket-sized field card listing bird taxa found in Hawai'i. \$.50 each. Call for postage price.

Hawaii Audubon Society logo t-shirts in white, natural, Pacific blue, jade, and gray, in sizes from S to XXL. Tank tops in white and grey, S to XL, same price. \$18.00 (\$15.00 plus \$3.00 postage and handling).

2004 Membership in Hawaii Audubon Society Regular US Member Foreign Membership (Airmail) (via bulk mail, not forwardable) \$ 15.00 Mexico \$ 21.00 First Class Mail\$ 21.00 Canada\$ 22.00 Junior Members (18 and under)\$ 10.00 All other countries \$ 28.00 Supporting Member\$100.00 Donations are tax deductible and gratefully accepted. Name Address City, State, Country, Zip _ Phone _ Email -Membership \$ ____ + Donation \$ _ = Total \$ _ New Membership Renewal Please make checks payable to Hawaii Audubon Society and mail to us at 850 Richards St., #505, Honolulu, HI 96813. PLEASE LET US KNOW IF YOUR ADDRESS CHANGES.



Nonprofit Organization
U.S. Postage
PAID

Honolulu, Hawaii Permit Number 1156

ADDRESS SERVICE REQUESTED

'ELEPAIO · 64:1 · FEBRUARY 2004

Calendar of Events

Table of Contents

March 15th, Monday, HAS Board Meeting:	
Open to all members, 6:30 to 8:30 p.m. at the HAS of	fice.
Education and Conservation Committees meet at 5:45	p.m.

February 1 to March12, HPU Art Gallery at Kane ohe campus presents "Birdwatching." See page 4.

February 16th, Monday, Program Meeting: (Sharon Reilly of Ducks Unlimited). See page 2.

Saturday, March 27th, Field Trip: Kahuku Dunes shoreline walk with Dr. Phil Bruner. See page 2.

Pacific and American Golden-Plovers: Reflections on Conservation Needs
February 16 Program Meeting
Results of Board Elections
Field Trips for 2004
Kolea Watch III-Spring 2004 2
Update on Studies of the Mid-Pacific Migration of Kolea Between Hawai'i and Alaska
Many Thanks 4
HPU Kane ohe Art Gallery Presents "Birdwatching" 4
Greenhouse Frog Invasion 4
HAS Products and Publications
Membership Application